

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Problem Image Mailbox.**



(11) (21) (C) **2,178,637**
(86) 1994/12/14
(87) 1995/06/22
(45) 1997/12/23

- (72) Pathak, Ram Dutta, GB
(72) Doughty, David George, GB
(73) SmithKline Beecham plc, GB
(51) Int. Cl.⁶ A61K 31/445, A61K 9/20
(30) 1993/12/15 (9325644.4) GB
(54) **COMPRIMES DE PAROXETINE ET LEUR PROCEDE DE
PREPARATION**
(54) **PAROXETINE TABLETS AND PROCESS TO PREPARE THEM**

SECTION 8 CORRECTION
SEE CERTIFICATE
CORRECTION - ARTICLES
VOIR CERTIFICATE

(57) L'invention concerne la paroxétine qui est préparée selon un procédé de formulation sans eau et qui est présentée sous forme de comprimés.

(57) Paroxetine which is formulated into tablets using a formulation process in which water is absent.





Bureau canadien

des brevets

Certificat de correction

Canadian Patent

Office

Certificate of Correction

Canadian Patent No. 2,178,637
Granted: December 23, 1997

Les corrections suivantes sont faites en
raison de l'article 8 de la *Loi sur les*
brevets et le document doit être lu tel
que corrigé.
In the Patent file and Patent grant:

1.

In claim 2, line 2 the words "followed by compression"
have been deleted.

The following corrections are made
pursuant to section 8 of the *Patent Act*
and the document should read as
corrected.


Agent certificateur / Certifying Officer

May 13, 1999
Date



Industrie
Canada

Industry
Canada

(CIPO 25)

Canada

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61K 31/445, 9/20	A1	(11) International Publication Number: WO 95/16448
		(43) International Publication Date: 22 June 1995 (22.06.95)

(21) International Application Number: PCT/EP94/04164

(22) International Filing Date: 14 December 1994 (14.12.94)

(30) Priority Data:
9325644.4 15 December 1993 (15.12.93) GB

(71) Applicant (for all designated States except US): SMITHKLINE
BEECHAM PLC [GB/GB]; New Horizons Court, Brentford,
Middlesex TW8 9EP (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PATHAK, Ram, Datta
[GB/GB]; SmithKline Beecham Pharmaceuticals, Great
Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ
(GB). DOUGHTY, David, George [GB/GB]; SmithKline
Beecham Pharmaceuticals, Mundella, Welwyn Garden City,
Hertfordshire AL7 1EY (GB).

(74) Agent: RUSSELL, Brian, J.; SmithKline Beecham, Corporate
Intellectual Property, SB House, Great West Road, Brent-
ford, Middlesex TW8 9BD (GB).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH,
CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG,
KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW,
NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT,
UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK,
ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI
patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE,
SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).

Published

With international search report.
Before the expiration of the time limit for amending the
claims and to be republished in the event of the receipt of
amendments.

2178637

(54) Title: PAROXETINE TABLETS AND PROCESS TO PREPARE THEM

(57) Abstract

Paroxetine which is formulated into tablets using a formulation process in which water is absent.

2 1 7 8 6 3 7

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:


1. A paroxetine formulation which is prepared on a commercial scale into tablets using a formulation process in which water is absent.
2. A formulation according to claim 1 in which the process is a dry direct compression of paroxetine followed by compression into tablets or a dry granulation of paroxetine followed by compression into tablets.
3. A formulation according to claim 1 or 2 in which the process for preparing it comprises the step of admixing paroxetine with dry excipients.
4. A formulation according to any one of claims 1 to 3 in which the paroxetine used in the process is in the form of the hydrochloride hemi-hydrate.

SECTION 5 CORRECTION
OF CERTIFICATE
CORRECTION - ARTICLE 8
OF CERTIFICATE

Paroxetine tablets and process to prepare them

The present invention relates to novel formulations and to the use of the formulation in the treatment and/or prevention of certain disorders.

- 5 US Patent 4,007,196 describes certain compounds which possess anti-depressant activity. One specific compound mentioned in this patent is known as paroxetine and which has the following formula:



10

This compound has been approved for human use and is being sold in many countries around the world as an anti-depressant agent.

It has been noticed that tablets of paroxetine often develop a pink hue which is highly undesirable.

- 15 To date, all tablets which have been sold have been formulated using an aqueous granulation process. It has surprisingly been found that formulation of paroxetine into tablets can be carried out reliably and on a commercial scale using a formulation process in which water is absent, such as by direct compression or by dry granulation.

20 It has also been surprisingly found that paroxetine formulated into a tablet using a process in which water is absent, is much less likely to develop a pink hue.

Accordingly, the present invention provides paroxetine which is formulated into tablets using a formulation process in which water is absent.

- 25 Examples of such a formulation process are dry direct compression of paroxetine or dry granulation of paroxetine followed by compression into tablets. The present invention therefore provides a formulation comprising direct compressed paroxetine admixed with dry excipients in the form of a tablet and a formulation comprising dry granulated and compressed paroxetine admixed with dry excipients in the form of a tablet.

30 It should be appreciated that the term "dry" means substantially "dry" as opposed to the wholesale addition of water which was previously employed in the wet granulation process.

Direct compression techniques are generally known in the art of pharmaceutical science. For example, paroxetine is conventionally admixed with dry excipients and compressed into tablets.

5 Dry granulation techniques are generally also known in the art of pharmaceutical science. For example, paroxetine is conventionally admixed with dry excipients and compressed into large slugs or roller compacted into ribbon-like strands. The compacted material is then suitably milled to produce a free flowing powder which is then compressed into tablets.

10 Additional excipients may then be added and mixed with the free flowing powder before being compressed into tablets.

Examples of excipients include calcium phosphate, microcrystalline cellulose, sodium starch glycollate and magnesium stearate which may be admixed in appropriate ratios.

15 It should be appreciated that particularly good results are obtained when microcrystalline cellulose is absent from the formulation, this is surprising as tablets formulated in the absence of microcrystalline cellulose are often prone to breaking up during manufacture or storage.

20 The paroxetine/excipient mixture may be compressed into an appropriate tablet shape. Preferred shapes include a pentagonal circumcircle, oval, round bi-convex or a tilt-tablet such as those described in US Patent 4,493,822.

Paroxetine when incorporated into the above-mentioned tablets is suitably, present as the hydrochloride hemi-hydrate form which may be prepared according to the procedures outlined in US Patent 4,721,723.

25 The amount of paroxetine present in the above-mentioned tablets is in the range of 10 to 100 mg of paroxetine as measured in terms of the "free base". Particularly preferred amounts include 10 mg, 20 mg, 30 mg, 40 mg and 50 mg of paroxetine as measured in terms of the "free base". Particularly preferred amounts include 20 mg, 30 mg and 40 mg of paroxetine as measured in terms of the "free base".

30 Suitable procedures for preparing paroxetine include those mentioned in US Patents 4,009,196, 4,902,801, 4,861,893 and 5,039,803 and PCT/GB 93/00721.

35 It has been mentioned that paroxetine has particular utility in the treatment of depression, paroxetine may also be used in the treatment of mixed anxiety and depression, obsessive compulsive disorders, panic, pain, obesity, senile dementia, migraine, bulimia, anorexia, social phobia and the depression arising from pre-menstrual tension and adolescence.

The present invention therefore also provides a method of treating or preventing any of the above disorders which comprises administering an effective or

prophylactic amount to a sufferer in need thereof of paroxetine which is formulated into a tablet using a process in which water is absent.

The present invention further provides a pharmaceutical composition comprising paroxetine which is formulated into a tablet using a process in which
5 water is absent for use in treating or preventing of the above disorders.

The present invention further provides the use of paroxetine which is formulated into a tablet using a process in which water is absent in the manufacture of a medicament for treating or preventing the above disorders.

The following examples illustrate the present invention:

10

Example 1

INGREDIENTS	20 mg Tablet	30mg Tablet
Paroxetine hydrochloride hemihydrate	22.67 mg	34.0 mg
Dicalcium Phosphate (DCP)	83.34 mg	125.0 mg
Microcrystalline Cellulose	50.67 mg	76.0 mg
Sodium Starch Glycollate	8.34 mg	12.5 mg
Magnesium Stearate	1.67 mg	2.5 mg
Tablet Weight	166.7 mg	250.0 mg

15 **Commercial source of the ingredients**

Dicalcium Phosphate Dihydrate	-	Emcompress or Ditab*
Microcrystalline Cellulose	-	Avicel PH 102*
Sodium Starch Glycollate	-	Explotab.*

20

* Trademarks

2178637

Method

1. Pass DCP through a screen and weigh it into a Planetary mixer.
- 5 2. Add 30 mesh Paroxetine to the bowl.
3. Add 20 mesh Avicel and Explotab and mix all the powders for 10 minutes.
4. Add magnesium Stearate and mix for 5 minutes.

10

Tablet into Pentagonal Tablets using the following punches:

	30 mg Tablet	9.5 mm	Circumcircle
15	20 mg Tablet	8.25 mm	Circumcircle

— The tablets are made satisfactorily on a single punch or a Rotary press.

Example 2

INGREDIENTS	10 mg Tablet	20 mg Tablet	30mg Tablet
Paroxetine hydrochloride hemihydrate	11.40 mg	22.80 mg	34.20 mg
Sodium Starch Glycollate	2.98 mg	5.95 mg	8.93 mg
Granular Dicalcium Phosphate (DITAB) or Dicafos	158.88 mg	317.75 mg	476.63 mg
Magnesium Stearate	1.75 mg	3.50 mg	5.25 mg
Tablet Weight	175.00 mg	350.00 mg	525.00 mg

5 Method

1. Paroxetine, Sodium Starch Glycollate and Dicalcium Phosphate Dihydrate are screened and mixed together in a suitable mixer.
(Planetary, Cuble or High Energy Shear mixer.)
2. Add Magnesium Stearate and compress it into a tablet using a single punch or Rotary Tablet machine.